



This plan should be completed by the student's personal diabetes health care team, including the parents/guardian. It should be reviewed with relevant school staff and copies should be kept in a place that can be accessed easily by the school nurse, trained diabetes personnel, and other authorized personnel.

| Date of Plan: | _ This plan is valid for the cu | urrent school year: | |
|-------------------------------|---------------------------------|---------------------|--|
| Student's Name: | Date of Birth: | | |
| Date of Diabetes Diagnosis: | type 1 | type 2 Other | |
| School: | School Phone Nur | nber: | |
| | Homeroom Teacher: | | |
| School Nurse: | Phone: | | |
| CONTACT INFORMATION | | | |
| M | | | |
| 0 | | | |
| ther/Guardian: | | | |
| Address: | | | |
| Telephone: Home | Work | _Cell: | |
| Email Address: | | | |
| Father/Guardian: | | | |
| | | | |
| Telephone: Home | Work | Cell: | |
| Email Address: | | | |
| S | | | |
| t | | | |
| udent's Physician/Health Care | e Provider: | | |
| Address: | | | |

| Telephone: | | |
|---------------------------|-------------------|-------|
| Email Address: | Emergency Number: | |
| Other Emergency Contacts: | | |
| Name: | Relationship: | |
| Telephone: Home | Work | Cell: |

CHECKING BLOOD GLUCOSE

| Target range of blood glucose: 70-130 mg/dL 70-180 mg/dL |
|--|
| Other: |
| Check blood glucose level: Before lunch Hours after lunch |
| ☐ 2 hours after a correction dose ☐ Mid-morning ☐ Before PE ☐ After |
| PE |
| ☐ Before dismissal ☐ Other: |
| ☐ As needed for signs/symptoms of low or high blood glucose☐ As needed for signs/symptoms of illness |
| Preferred site of testing: |
| Brand/Model of blood glucose meter: |
| Note: The fingertip should always be used to check blood glucose level if hypoglycemia is suspected. |
| Student's self-care blood glucose checking skills: |
| ☐ Independently checks own blood glucose |
| May check blood glucose with supervision |
| Requires school nurse or trained diabetes personnel to check blood glucose |
| Continuous Glucose Monitor (CGM): Yes No Brand/Model: Alarms set for: (low) and (high) |
| Note: Confirm CGM results with blood glucose meter check before taking action on sensor blood glucose level. If student has symptoms or signs of hypoglycemia, check fingertip blood glucose level regardless of CGM |
| HYPOGLYCEMIA TREATMENT |
| |
| Student's usual symptoms of hypoglycemia (list below): |
| If exhibiting symptoms of hypoglycemia, OR if blood glucose level is less than mg/dL, give a quick-acting glucose product equal to grams of carbohydrate. |

| Recheck blood glucose in 10-15 minutes and repeat treatment if blood glucose level is |
|---|
| less than mg/dL. |
| Additional treatment: |

HYPOGLYCEMIA TREATMENT (Continued)

| Follow physical activity and sports orders (see page 7). |
|---|
| • If the student is unable to eat or drink, is unconscious or unresponsive, or is having |
| seizure activity or convulsions (jerking movements), give: |
| • Glucagon: |
| • Site for glucagon injection: arm thigh Other: |
| • Call 911 (Emergency Medical Services) and the student's parents/guardian. |
| Contact student's health care provider. |
| |
| HYPERGLYCEMIA TREATMENT |
| |
| |
| |
| |
| Student's usual symptoms of hyperglycemia (list below): |
| |
| |
| Check Urine Blood for ketones every hours when blood glucose levels |
| are abovemg/dL. |
| For blood glucose greater thanmg/dL AND at least hours since last insulin dose, give correction dose of insulin (see orders below). |
| For insulin pump users: see additional information for student with insulin pump. |
| Give extra water and/or non-sugar-containing drinks (not fruit juices):ounces per hour. |
| Additional treatment for ketones: |
| Follow physical activity and |
| sports orders (see page 7). |
| • Notify namenta/ayandian of anget of hymanalysassis |

- Notify parents/guardian of onset of hyperglycemia.
- If the student has symptoms of a hyperglycemia emergency, including dry mouth, extreme thirst, nausea and vomiting, severe abdominal pain, heavy breathing or shortness of breath, chest pain, increasing sleepiness or lethargy, or depressed level of consciousness: Call 911 (Emergency Medical Services) and the student's parents/guardian.
- Contact student's health care provider.

| Diabetes Medical Management Plan (DMMP) — page 4 |
|---|
| INSULIN THERAPY |
| Insulin delivery device: syringe insulin pen insulin pump |
| Type of insulin therapy at school: Adjustable Insulin Therapy Fixed Insulin Therapy No insulin |
| Adjustable Insulin Therapy |
| • Carbohydrate Coverage/Correction Dose: |
| N |
| a |
| me of insulin: |
| • Carbohydrate Coverage: |
| Insulin-to-Carbohydrate Ratio: |
| Lunch: 1 unit of insulin per grams of carbohydrate |
| Snack: 1 unit of insulin per grams of carbohydrate |
| |
| Carbohydrate Dose Calculation Example |
| Grams of carbohydrate in meal |
| Insulin-to-carbohydrate ratio = units of insulin |
| |
| • Correction Dose: |
| Correction Dose Calculation Example |
| Actual Blood Clucosa Target Blood Clucosa |
| Blood Glucose Correction Factor/Insulin Sensitivity Factor =units of insulin |
| |
| Blood Glucose Correction Factor/Insulin Sensitivity Factor = |
| Target blood glucose = mg/dL |
| Correction dose scale (use instead of calculation above to determine insulin correction dose): |
| Blood glucose to mg/dL give units |
| Blood glucose tomg/dL give units |
| Blood glucose to mg/dL give units |
| Blood glucose to mg/dL give units |

INSULIN THERAPY (Continued)

| When to give insulin: |
|---|
| Lunch Carbahydrata agyaraga anly |
| Carbohydrate coverage only |
| Carbohydrate coverage plus correction dose when blood glucose is greater than mg/dL and hours since last insulin dose. |
| Other: nours since last insumit dose. |
| Snack |
| ☐ No coverage for snack |
| Carbohydrate coverage only |
| Carbohydrate coverage plus correction dose when blood glucose is greater than |
| mg/dL and hours since last insulin dose. |
| Other: |
| |
| Correction dose only: |
| For blood glucose greater than mg/dL AND at leasthours since last |
| insulin dose. |
| Other: |
| |
| Fixed Insulin Therapy |
| N |
| a |
| me of insulin: |
| Units of insulin given pre-lunch daily |
| Units of insulin given pre-snack daily |
| Other: |
| Parental Authorization to Adjust Insulin Dose: |
| |
| Yes No Parents/guardian authorization should be obtained before administering a correction dose. |
| Yes No Parents/guardian are authorized to increase or decrease correction dose scale within the following range: +/- units of insulin. |
| Yes No Parents/guardian are authorized to increase or decrease insulin-to-carbohydrate ratio within the following range: _ units per prescribed grams of carbohydrate, +/- grams of carbohydrate. |
| Yes No Parents/guardian are authorized to increase or decrease fixed insulin |

dose within the following range: +/- _ units of insulin.

INSULIN THERAPY (Continued)

| Student's self-care insulin administration skill | |
|---|-------------------------------------|
| Yes No Independently calculates and gives | s own injections |
| ☐ Yes ☐ No May calculate/give own injections | with supervision |
| Yes No Requires school nurse or trained dinjections | iabetes personnel to calculate/give |
| ADDITIONAL INFORMATION FOR STUDENT | WITH INSULIN PUMP |
| Brand/Model of pump: Typ | oe of insulin in pump: |
| Basal rates during school: | |
| Type of infusion set: | |
| For blood glucose greater than mg/dLhours after correction, consider pump parents/guardian. | |
| For infusion site failure: Insert new infusion set | and/or replace reservoir. |
| For suspected pump failure: suspend or remove pen. | pump and give insulin by syringe or |
| Physical Activity | |
| May disconnect from pump for sports activities | |
| Set a temporary basal rate Yes No———Suspend pump use Yes No | - % temporary basal for hours |
| Student's self-care pump skills: | Independent? |
| Count carbohydrates | ☐ Yes ☐ No |
| Bolus correct amount for carbohydrates consumed | ☐ Yes ☐ No |
| Calculate and administer correction bolus | ☐ Yes ☐ No |
| Calculate and set basal profiles | ☐ Yes ☐ No |
| Calculate and set temporary basal rate | ☐ Yes ☐ No |
| Change batteries | ☐ Yes ☐ No |
| Disconnect pump | ☐ Yes ☐ No |
| Reconnect pump to infusion set | ☐ Yes ☐ No |
| Prepare reservoir and tubing | ☐ Yes ☐ No |
| Insert infusion set | ☐ Yes ☐ No |
| Troubleshoot alarms and malfunctions | ☐ Yes ☐ No |

| OTHER DIABETES N | MEDICATIONS | | | |
|---|---------------------------------|-----------|----------------------|----------------------|
| Name: | Dose:Dose: | | Route: | Times given: |
| Name: | Dose: | | Route: | Times given: |
| MEAL PLAN | | | | |
| Meal/Snack | Time | Car | bohydrate Conte | nt (grams) |
| Breakfast | | | | , |
| Mid-morning snack | | | | |
| Lunch | | | to | |
| Mid-afternoon snack | | | to | |
| Other times to give snae | cks and content/an | nount: | | |
| Instructions for when for sampling event): | ood is provided to | | | class party or food |
| Special event/party food | d permitted: 🔲 F | Parents/g | guardian discretion | |
| | | student d | iscretion | |
| Student's self-care nu Yes No Inde | | carbohy | lrates | |
| ☐ Yes ☐ No May | count carbohydra | tes with | supervision | |
| - | uires school nurse/ hydrates | trained (| diabetes personnel | to count |
| PHYSICAL ACTIVIT | TY AND SPORTS | S | | |
| A quick-acting source of sugar-containing juice sports. | _ | _ | | |
| Student should eat 1 | 5 grams | rams of | carbohydrate 🔲 | other |
| ☐ before ☐ every | 30 minutes during | g 🔲 a | fter vigorous physi | ical activity |
| other | | | | • |
| If most recent blood gluphysical activity when | cose is less than _ | 4 | _mg/dL, student c | |
| Avoid physical activity blood ketones are mode | _ | se is gre | eater than | _ mg/dL or if urine/ |
| (Additional information | n for student on ins | sulin pui | np is in the insulin | section on page 6.) |

DISASTER PLAN

| To prepare for an unplanned disaster or emosupply kit from parent/guardian. | ergency (72 HOURS), obtain emergency |
|---|--|
| Continue to follow orders contained in | this DMMP. |
| Additional insulin orders as follows: | |
| Other: | |
| SIGNATURES | |
| This Diabetes Medical Management Plan ha | as been approved by: |
| Student's Physician/Health Care Provider | Date |
| I, (parent/guardian:) | give permission to the school nurse |
| or another qualified health care professiona | l or trained diabetes personnel of |
| (school:) | to perform and carry out the diabetes care |
| tasks as outlined in (student:) | 's Diabetes Medical Management |
| Plan. I also consent to the release of the info | ormation contained in this Diabetes Medical |
| Management Plan to all school staff member | ers and other adults who have responsibility |
| for my child and who may need to know thi | s information to maintain my child's health |
| and safety. I also give permission to the sch | ool nurse or another qualified health care |
| professional to contact my child's physician | /health care provider. |
| | |
| | |
| | |
| | |
| | Date |
| Acknowledged and received by: | Date |
| Student's Parent/Guardian | Date |

| Stu | dent's | Parent/ | Guardiar | 1 |
|-----|--------|---------|----------|---|
| | | 1 (11) | | |

School Nurse/Other Qualified Health Care Personnel